

FINITENESS OF ARITHMETIC HYPERBOLIC REFLECTION GROUPS

IAN AGOL, MIKHAIL BELOLIPETSKY, PETER STORM, AND KEVIN WHYTE

ABSTRACT. We prove that there are only finitely many conjugacy classes of arithmetic maximal hyperbolic reflection groups.

1. INTRODUCTION

A hyperbolic reflection group Γ is a discrete group generated by reflections in the faces of a hyperbolic polyhedron $P \subset \mathbb{H}^n$. We may assume that the dihedral angles of P are of the form π/n , $n \geq 2$, in which case P forms a fundamental domain for the action of Γ on $\mathbb{H}^n$. If P has finite volume, then $\mathbb{H}^n/\Gamma = \mathcal{O}$ is a hyperbolic orbifold of finite volume, which is obtained by “mirroring” the faces of P . A reflection group Γ is maximal if there is no reflection group Γ' such that $\Gamma < \Gamma'$. We defer the definition of arithmetic groups until later, but a theorem of Margulis implies that Γ is arithmetic if and only if $[\text{Comm}(\Gamma) : \Gamma] = \infty$, where $\text{Comm}(\Gamma) = \{g \in \text{Isom}(\mathbb{H}^n) \mid [\Gamma : g^{-1}\Gamma g \cap \Gamma] < \infty\}$ [23]. The main theorem of this paper is that up to conjugacy in $\text{Isom}(\mathbb{H}^n)$ there are only finitely many arithmetic groups which are maximal hyperbolic reflection groups. If one does not assume that the groups are maximal, then there is no corresponding finiteness result. For example, using a method of [33] one can see that there are all right angle reflection groups which then contain infinitely many finite index reflection subgroups, by a sequence of doublings along faces. These examples are known up to dimension 8, but recently examples of infinite families of arithmetic reflection groups up to dimension 19 have been given by Allcock [2].

Finiteness results of this sort were proven before by Nikulin [24, 25, 26] for dimension $n \geq 10$. Subsequently Vinberg showed that there are no uniform (*i.e.* cocompact) reflection groups in dimension $n \geq 30$ [40]. Our results are independent of their work, in the sense that we obtain a finiteness result in each dimension and then appeal to the result of Prokhorov that reflection groups do not exist in high enough dimensions [30]. The argument of this

Date: December 4, 2006.

2000 *Mathematics Subject Classification.* 20F55, 22E40, 11F06, 11H56.

Agol partially supported by NSF grant DMS-0504975 and the Guggenheim Foundation.

Belolipetsky thanks MPIM in Bonn for hospitality and support during the summer of 2006.

Storm partially supported by NSF grant DMS-0603711.

Whyte partially funded by NSF Career grant DMS-0349290 and the Sloan Foundation.

paper generalizes an argument of Long-MacLachlan-Reid [21], which implies that there are only finitely many arithmetic minimal (or congruence) hyperbolic 2-orbifolds with bounded genus. Their argument is in fact a generalization of an argument of Zograf [43], who reproved that there are only finitely many congruence groups Γ commensurable with $\mathrm{PSL}(2, \mathbb{Z})$ such that $\mathbb{H}^2/\Gamma$ has genus 0 (This was proven originally by Dennin [9, 10], and was known as Rademacher's conjecture). These arguments were generalized in [1] to show that there are only finitely many maximal arithmetic reflection groups in dimension 3. Thus, our result is only new for cocompact maximal arithmetic groups in dimensions $4 \leq n \leq 9$. One key ingredient of these arguments is a theorem of Burger and Sarnak [8], which implies that there is a lower bound on the first eigenvalue of a congruence arithmetic hyperbolic group defined by a quadratic form. The other key ingredient is an inequality of Li-Yau [20], generalized to orbifolds, which allows us to give an upper bound on the first eigenvalue of a maximal quotient of an arithmetic reflection group. One of the new ingredients of the present work is the use of some results on the distribution of covolumes of maximal arithmetic subgroups obtained in [3].

Remark: As this manuscript was being prepared, a short preprint appeared by Nikulin proving the finiteness of arithmetic maximal reflection groups in all dimensions [27]. Nikulin's proof makes use of Nikulin's previous results [24, 25], as well as the finiteness results for maximal arithmetic reflection groups in dimensions two and three [21, 1]. The methods in this paper are completely independent of Nikulin's methods. A final classification of maximal arithmetic reflection groups might combine the techniques of both approaches, so it seemed interesting to publish our result, even with the appearance of Nikulin's preprint.

2. CONFORMAL VOLUME OF ORBIFOLDS

Conformal volume was first defined by Li-Yau [20], partially motivated by generalizing results on surfaces due to Yang-Yau [42], Hersch [15], and Szegő [35]. We generalize this notion to orbifolds. Let $(\mathcal{O}, g)$ be a complete Riemannian orbifold, possibly with boundary. Let $|\mathcal{O}|$ denote the underlying topological space. Denote the volume form by dv_g , and the volume by $\mathrm{Vol}(\mathcal{O}, g)$. Let $\mathrm{Möb}(\mathbb{S}^n)$ denote the conformal transformations of $\mathbb{S}^n$. It is well-known that $\mathrm{Möb}(\mathbb{S}^n) = \mathrm{Isom}(\mathbb{H}^{n+1})$. The topological space $|\mathcal{O}|$ has a dense open subset which is a Riemannian manifold. We will say that a map $\varphi : |\mathcal{O}_1| \rightarrow |\mathcal{O}_2|$ is *PC* if it is a continuous map which is piecewise a conformal immersion. Clearly, if $\varphi : |\mathcal{O}| \rightarrow \mathbb{S}^n$ is PC, and $\mu \in \mathrm{Möb}(\mathbb{S}^n)$, then $\mu \circ \varphi$ is also PC. Let $(\mathbb{S}^n, \mathrm{can})$ be the canonical round metric on the n -sphere.

Definition 2.1. For a piecewise smooth map $\varphi : |\mathcal{O}| \rightarrow (\mathbb{S}^n, \mathrm{can})$, define

$$V_{PC}(n, \varphi) = \sup_{\mu \in \mathrm{Möb}(\mathbb{S}^n)} \mathrm{Vol}(\mathcal{O}, (\mu \circ \varphi)^*(\mathrm{can})).$$

If there exists a PC map $\varphi : |\mathcal{O}| \rightarrow \mathbb{S}^n$, then we also define

$$V_{PC}(n, \mathcal{O}) = \inf_{\varphi : |\mathcal{O}| \rightarrow \mathbb{S}^n_{PC}} V_{PC}(n, \varphi).$$

$V_{PC}(n, \mathcal{O})$ is denoted the (n -dimensional) *piecewise conformal volume* of $\mathcal{O}$.

Remark: It seems likely that our definition of piecewise conformal volume coincides with conformal volume of Li-Yau for manifolds, but we have not checked this (it would suffice to show that a PC map can be approximated by conformal maps).

If there exists a piecewise isometric map $\varphi : (|\mathcal{O}|, g) \rightarrow \mathbb{E}^n$ for some n , then clearly $V_{PC}(n, \mathcal{O})$ is well-defined, since $\mathbb{E}^n$ has a conformal embedding into $\mathbb{S}^n$. Let $\Sigma \subset \mathcal{O}$ be the singular locus of $\mathcal{O}$. By the Nash embedding Theorem [14], there is an isometry $\rho : \mathcal{O} \setminus \Sigma \rightarrow \mathbb{E}^N$, for N large enough. Then ρ extends continuously to an isometric map $\bar{\rho} : \mathcal{O} \rightarrow \mathbb{E}^N$. Thus, $V_{PC}(N, \mathcal{O})$ is well-defined. We record some basic facts about conformal volume, generalizing some of the facts in [20, §1] which carry over to our notion of conformal volume for orbifolds.

Fact 1: If $\mathcal{O}$ admits a PC map ψ of geometric degree d onto another orbifold $\mathcal{P}$, then

$$V_{PC}(n, \mathcal{O}) \leq |d| V_{PC}(n, \mathcal{P}).$$

This follows because for any PC map $\varphi : \mathcal{P} \rightarrow \mathbb{S}^n$, the composition $\varphi \circ \psi : \mathcal{O} \rightarrow \mathbb{S}^n$ satisfies $V_{PC}(n, \varphi \circ \psi) \leq d V_{PC}(n, \varphi)$.

Fact 2: Since the embedding $\mathbb{S}^n \hookrightarrow \mathbb{S}^{n+1}$ is an isometry, it's clear that $V_{PC}(n, \mathcal{O}) \geq V_{PC}(n+1, \mathcal{O})$. Define the piecewise conformal volume $V_{PC}(\mathcal{O}) = \lim_{n \rightarrow \infty} V_{PC}(n, \mathcal{O})$.

Fact 3: If $\mathcal{O}$ is of dimension m , and $\varphi : |\mathcal{O}| \rightarrow \mathbb{S}^n$ is a PC map, then

$$V_{PC}(n, \varphi) \geq V_{PC}(n, \mathbb{S}^m) = \text{Vol}(\mathbb{S}^m).$$

The same argument as in Li-Yau works here: “blow up” about a smooth manifold point of $\varphi(|\mathcal{O}|)$ so that the image converges in the Hausdorff topology to a geodesic sphere of dimension m . More precisely, let $x \in \mathcal{O}$ be a manifold point of $\mathcal{O}$ such that the differential $d\varphi_x$ has rank m . Let $\mu_t : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ be a Möbius transformation preserving $\mathbb{S}^n$, fixing $\pm\varphi(x)$, and translating the origin $0 \in \mathbb{R}^{n+1}$ to $-t\varphi(x)$, $0 \leq t < 1$. Let $U \subset \mathcal{O}$ be a small neighborhood of x such that $rk(d\varphi_y) = m$, for all $y \in U$. Then $\mu_t(\varphi(U))$ converges to the geodesic sphere of dimension m going through $\pm\varphi(x)$, and tangent to $d\varphi_x(T_x\mathcal{O})$ as $t \rightarrow 1$. Thus, we see that $V_{PC}(n, \varphi) \geq \text{Vol}(\mathbb{S}^m)$.

Fact 4: If $\mathcal{O}$ is an embedded suborbifold of the orbifold $\mathcal{P}$, and $\varphi : |\mathcal{P}| \rightarrow \mathbb{S}^n$ is PC, then $V_{PC}(n, \varphi) \geq V_{PC}(n, \varphi|_{|\mathcal{O}|})$. Thus, $V_{PC}(n, \mathcal{P}) \geq V_{PC}(n, \mathcal{O})$.

For example, suppose $\iota : P \subset \mathbb{S}^n$ is an n -dimensional submanifold, with the conformal structure induced from the embedding. Then $V_{PC}(n, \iota) \leq V_{PC}(\mathbb{S}^n)$, by fact 4. But by fact 3, $V_{PC}(n, \iota) \geq V_{PC}(\mathbb{S}^n)$, and therefore we have shown that $V_{PC}(n, \iota) = V_{PC}(P) = V_{PC}(\mathbb{S}^n) = \text{Vol}(\mathbb{S}^n)$.

Fact 5: If $\mathcal{O} = \mathcal{O}_1 \cup \mathcal{O}_2$, and $\varphi : \mathcal{O} \rightarrow \mathbb{S}^n$, then

$$V_{PC}(\mathcal{O}, \varphi) \leq V_{PC}(\mathcal{O}_1, \varphi|_{\mathcal{O}_1}) + V_{PC}(\mathcal{O}_2, \varphi|_{\mathcal{O}_2}).$$

3. FINITE SUBGROUPS OF $O(n+1)$

In this section, we give a bound on the conformal volume of $\mathbb{S}^n/\Gamma$, where $\Gamma < O(n+1)$ is a finite subgroup. For $\Gamma < O(n+1)$ finite, define the orbifold $M_\Gamma^n = \mathbb{S}^n/\Gamma$.

Theorem 3.1. *There is a function $C(n)$ such that if $\Gamma < O(n+1)$, then*

$$V_{PC}(M_\Gamma^n) < C(n)|\Gamma|^{2n}.$$

Proof. First, we need a lemma.

Lemma 3.2. *If we have k hyperplanes in $\mathbb{R}^n$ going through the origin, where $k > 0, n > 0$, then they separate $\mathbb{R}^n$ into at most $f(k, n) = 2 \sum_{i=0}^{n-1} \binom{k-1}{i} = O(k^{n-1})$ regions.*

Proof. We may assume the hyperplanes are in general position. The proof is by lexicographic induction on (k, n) . Clearly, $f(k, 1) = 2$, $f(1, n) = 2$. Consider k hyperplanes $\{P_1, \dots, P_k\}$, $P_i \subset \mathbb{R}^n$, $k > 1, n > 1$. Then the hyperplanes $\{P_1, \dots, P_{k-1}\}$ separate $\mathbb{R}^n$ into at most $f(k-1, n)$ regions, by the inductive hypothesis. Also, $\{P_1 \cap P_k, \dots, P_{k-1} \cap P_k\}$ is a collection of $k-1$ hyperplanes in $P_k \cong \mathbb{R}^{n-1}$, and thus they divide P_k into at most $f(k-1, n-1)$ regions, by the inductive hypothesis. Each region of $\mathbb{R}^n \setminus (P_1 \cup \dots \cup P_{k-1})$ which meets P_k will be divided into two, and there are at most $f(k-1, n-1)$ such regions. The regions which don't meet P_k will remain unchanged. Thus, $|\mathbb{R}^n \setminus (P_1 \cup \dots \cup P_k)| \leq f(k-1, n) + f(k-1, n-1) = f(k, n)$. This follows since

$$\begin{aligned} f(k-1, n) + f(k-1, n-1) &= 2 \sum_{i=0}^{n-1} \binom{k-2}{i} + 2 \sum_{i=0}^{n-2} \binom{k-2}{i} \\ &= 2 \sum_{i=0}^{n-1} \binom{k-2}{i} + \binom{k-2}{n-1} = 2 \sum_{i=0}^{n-1} \binom{k-1}{i} = f(k, n). \end{aligned}$$

□

We will actually find a piecewise isometric map $f : \mathbb{S}^n/\Gamma \rightarrow \mathbb{S}^N$ for some N , in which the number of isometrically embedded pieces is bounded by a polynomial in $|\Gamma| = m$. This will immediately imply the result, using facts 3 through 5 about conformal volume.

Define the orbi-covering $\pi_\Gamma : \mathbb{S}^n \rightarrow M_\Gamma^n$. Let $\Gamma = \{\gamma_1, \gamma_2, \dots, \gamma_m\}$. For each element $\gamma \in \Gamma$, right multiplication by γ induces a permutation of $\{\gamma_1, \gamma_2, \dots, \gamma_m\}$, which we identify with a permutation of $\{1, 2, \dots, m\}$. Notice this does not give an action of Γ , since composing right multiplications

reverses the order of group multiplication. Use these permutations to define, for each $\gamma \in \Gamma$, a permutation of the coordinates of $\prod_{i=1}^m \overline{\mathbb{B}}^{n+1}$ given by

$$\gamma \cdot (x_1, \dots, x_m) = (x_{\gamma \cdot 1}, \dots, x_{\gamma \cdot m}).$$

Define $N = m(n+1) - 1$ and

$$\begin{aligned} \Psi : \mathbb{S}^n &\longrightarrow \mathbb{S}^N \subset \prod_{i=1}^m \overline{\mathbb{B}}^{n+1} \\ x &\longmapsto m^{-\frac{1}{2}}(\gamma_1 x, \gamma_2 x, \dots, \gamma_m x). \end{aligned}$$

$\mathbb{S}^n$ is taken by Ψ to a totally geodesic submanifold of $\mathbb{S}^N$, since it is (up to rotation in $O(N+1)$) the restriction of the diagonal map $\mathbb{R}^{n+1} \rightarrow \mathbb{R}^{N+1}$, rescaled to be an isometry. For each element $\gamma \in \Gamma$ the map Ψ is γ -equivariant, because

$$\Psi(\gamma \cdot x) = m^{-\frac{1}{2}}(\gamma_1 \cdot \gamma \cdot x, \dots, \gamma_m \cdot \gamma \cdot x) = m^{-\frac{1}{2}}(\gamma_{\gamma \cdot 1} x, \dots, \gamma_{\gamma \cdot m} x) = \gamma \cdot \Psi(x)$$

for any $x \in \mathbb{S}^n$.

Let Σ_m be the permutation group on m elements. The map Ψ therefore descends to a map

$$\Upsilon : M_\Gamma^n \longrightarrow \mathbb{S}^N / \Sigma_m = Y.$$

The action of Σ_m on $\mathbb{S}^N$ induces an embedding $\Sigma_m < O(N+1)$. By permuting the coordinates of $\mathbb{R}^{N+1}$, the symmetric group $\Sigma = \Sigma_{N+1}$ acts isometrically as a reflection group on $\mathbb{S}^N$, such that $\Sigma_m < \Sigma$. Similar to what we did above, define the orbi-covering $\pi_\Sigma : Y \longrightarrow M_\Sigma^N$ and an isometric embedding $\iota_\Sigma : |M_\Sigma^N| \longrightarrow \mathbb{S}^N$, which exists since we may identify $|M_\Sigma^N|$ with the Coxeter polyhedron for Σ . Let $\{S_k\}_{k \in K}$ be the collection of $\binom{N+1}{2}$ hyperspheres given by the fixed point sets of the reflections in Σ .

The goal is now to estimate $V_{PC}(M_\Gamma^n, (\iota_\Sigma \circ \pi_\Sigma \circ \Upsilon))$ from above.

Consider how $\Psi(\mathbb{S}^n)$ is cut up by the collection $\{S_k\}_{k \in K}$. It is cut up into at most $f\left(\binom{N+1}{2}, n+1\right)$ convex sub-polyhedra $\{C_\sigma, \sigma \in \Sigma\}$ by the planes $\{\Psi(\mathbb{S}^n) \cap S_k\}_{k \in K}$ (For most $\sigma \in \Sigma$, $C_\sigma = \emptyset$). Then each C_σ is isometrically and totally geodesically projected to $\mathbb{S}^N$ via the map $\iota_\Sigma \circ \pi_\Sigma \circ \Upsilon$. Postcomposition by a conformal automorphism of $\mathbb{S}^N$ can then make the volume of the image of C_σ no more than the volume of a unit n -sphere. By the combination of facts 3 through 5 of section 2, we can conclude that

$$V_{PC}(M_\Gamma^n, (\iota_\Sigma \circ \pi_\Sigma \circ \Upsilon)) \leq f\left(\binom{m(n+1)}{2}, n+1\right) \cdot \text{Vol}(\mathbb{S}^n) \leq C(n)|\Gamma|^{2n},$$

for some constant $C(n)$. $\square$

Remark: It turns out that a finite subgroup $\Gamma < O(n+1)$ does not necessarily embed in a finite reflection subgroup of $O(n+1)$. Thus, the strategy of [1] does not generalize to higher dimensions. It is an interesting problem to try to get better estimates on $V_{PC}(M_\Gamma^n)$. The method we have used to estimate this quantity is probably far from optimal. In fact, in

the above theorem we have constructed maps which potentially have very high multiplicity, which is exactly the sort of thing one wants to avoid in estimating conformal volume.

4. EIGENVALUE BOUNDS

In this section, we observe that the argument of [11, Thm. 2.2] generalizes to our context of orbifolds (their theorem sharpens Corollary 3 from section 2 of [20]). If $(\mathcal{O}, g)$ is a Riemannian orbifold, then $\lambda_1(\mathcal{O}, g)$ is the first non-zero eigenvalue of the operator $-\tilde{\Delta}_g$ on $\mathcal{O}$, where $\tilde{\Delta}_g$ is the unique self-adjoint extension of the Laplacian on $\mathcal{O}$. We will be using the variational characterization of $\lambda_1(\mathcal{O}, g)$ in terms of the Rayleigh-Ritz quotient.

Lemma 4.1. *Let $\mathcal{O}$ be an orbifold of finite volume. Let $f \in W^{1,2}(\mathcal{O})$ be an L^2 function with L^2 distributional derivative, and such that $\int_{\mathcal{O}} f dv_g = 0$. Then*

$$\lambda_1(\mathcal{O}, g) \int_{\mathcal{O}} f^2 dv_g \leq \int_{\mathcal{O}} |\nabla f|^2 dv_g.$$

Proof. Let A be a self-adjoint operator on a Hilbert space $\mathcal{H}$, such that A is semi-bounded below. This means that there exists a constant k , such that for all $\psi \in \mathcal{H}$, $\langle \psi, A\psi \rangle \geq k \langle \psi, \psi \rangle$. The Laplacian Δ with domain restricted to compactly supported, C^∞ functions on $\mathcal{O}$ is essentially self-adjoint ([12, Thm. 1.9]), with self-adjoint extension $\tilde{\Delta}$. In the case of the self-adjoint operator $-\tilde{\Delta}$, we have $-\tilde{\Delta}$ is semi-bounded below, with constant $k = 0$. This follows from the fact that $C_c^\infty(\mathcal{O})$ is dense in $\text{Domain}(\tilde{\Delta})$, and for $f \in C_c^\infty(\mathcal{O})$ integration by parts gives $\int_{\mathcal{O}} -f \tilde{\Delta} f dv_g = \int_{\mathcal{O}} |\nabla f|^2 dv_g$, which is non-negative. We have $\lambda_0(-\tilde{\Delta}) = 0$, with constant eigenfunction. Therefore, the Rayleigh-Ritz principle for semi-bounded below operators [31, XIII] gives

$$\lambda_1(\mathcal{O}, g) = \inf \left\{ \frac{\int_{\mathcal{O}} -f \tilde{\Delta} f dv_g}{\int_{\mathcal{O}} f^2 dv_g} \mid f \in \text{Domain}(\tilde{\Delta}), \int_{\mathcal{O}} f dv_g = 0 \right\}.$$

Since $C_c^\infty(\mathcal{O})$ is dense in $\text{Domain}(\tilde{\Delta})$, we may take the above infimum over functions $f \in C_c^\infty(\mathcal{O})$. Then integrating by parts, we have

$$\lambda_1(\mathcal{O}, g) = \inf \left\{ \frac{\int_{\mathcal{O}} |\nabla f|^2 dv_g}{\int_{\mathcal{O}} f^2 dv_g} \mid f \in C_c^\infty(\mathcal{O}), \int_{\mathcal{O}} f dv_g = 0 \right\}.$$

Then we use again the fact that $C_c^\infty(\mathcal{O})$ is dense in the Sobolev space $W^{1,2}(\mathcal{O})$ to see that we may take the above infimum over functions in $W^{1,2}(\mathcal{O})$. $\square$

Theorem 4.2. *(see [11, Thm. 2.2]) Let $(\mathcal{O}, g)$ be a Riemannian orbifold of dimension m of finite volume. If $\varphi : |\mathcal{O}| \rightarrow \mathbb{S}^n$ is a PC map,*

$$\lambda_1(\mathcal{O}, g) \text{Vol}(\mathcal{O}, g)^{\frac{2}{m}} \leq m V_{PC}(n, \varphi)^{\frac{2}{m}}.$$

Proof. Let $\mathbf{X} = (X_1, \dots, X_{n+1})$ be the coordinate functions on $\mathbb{R}^{n+1}$. Then $\sum_{i=1}^{n+1} X_i^2 = 1$, restricted to the unit sphere $\mathbb{S}^n \subset \mathbb{R}^{n+1}$.

Lemma 4.3. *There exists $\mu \in \text{Möb}(\mathbb{S}^n)$ such that $\int_{\mathcal{O}} \mathbf{X} \circ \mu \circ \varphi \, dv_g = \mathbf{0}$.*

Proof. Let $\mathbf{x} \in \mathbb{S}^n$. For $0 \leq t < 1$, $t\mathbf{x} \in \mathbb{B}^{n+1}$ (the open unit ball), let $\mu_{t\mathbf{x}} \in \text{Möb}(\mathbb{S}^n) = \text{Isom}(\mathbb{B}^{n+1})$ be the hyperbolic translation along the ray $\mathbb{R}\mathbf{x}$ taking $\mathbf{0}$ to $t\mathbf{x}$ (thus, $\mu_{0\mathbf{x}} = \mu_{\mathbf{0}} = \text{Id}$). Let $H(t\mathbf{x}) = \frac{1}{\text{Vol}(\mathcal{O}, g)} \int_{\mathcal{O}} \mathbf{X} \circ \mu_{t\mathbf{x}} \circ \varphi \, dv_g$. We may think of H as defining a continuous function $H : \mathbb{B}^{n+1} \rightarrow \mathbb{B}^{n+1}$, which gives the center of mass of the measure coming from $\varphi_* dv_g$, where we take the point $-t\mathbf{x}$ to the origin of the sphere $\mathbb{S}^n = \partial\mathbb{B}^{n+1}$ by the conformal map $\mu_{t\mathbf{x}}$. Let $x_i \in \mathbb{B}^{n+1}$ be a sequence of points approaching $x_\infty \in \mathbb{S}^n$. We want to show that H extends to a map fixing $\partial\mathbb{B}^{n+1}$. To see this, let $U_\delta = \mathbb{S}^n \setminus B_\delta(-x_\infty)$ be a neighborhood of x_∞ . Then the fact that φ is PC implies that for any $\epsilon > 0$, there exists $\delta > 0$, such that $\text{Vol}(\varphi^{-1}(U_\delta)) > \text{Vol}(\mathcal{O})(1 - \epsilon)$. Then we may find $N > 0$, such that $\mu_{x_i}(U_\delta) \subset B_\epsilon(x_\infty)$, for $i > N$. Thus, we may see that $H(x_i) \in B_\epsilon(x_\infty)$. So $\lim_{i \rightarrow \infty} H(x_i) = x_\infty$. So we see that H extends to a continuous function $\overline{H} : \overline{\mathbb{B}}^{n+1} \rightarrow \overline{\mathbb{B}}^{n+1}$ such that $\overline{H}|_{\mathbb{S}^n} = \text{Id}|_{\mathbb{S}^n}$, and $\overline{H}|_{\mathbb{B}^{n+1}} = H$. Thus, H is onto, so there exists $\mathbf{y} \in \mathbb{B}^{n+1}$ such that $H(\mathbf{y}) = \mathbf{0}$, and we take $\mu = \mu_{\mathbf{y}}$. $\square$

Now, replace φ with $\mu \circ \varphi$, noting that this is still a PC map. Then $X_i \circ \varphi$ may be used as test functions in the Rayleigh quotient, since they are L^2 orthogonal to the constant function, and lie in the Sobolev space $W^{1,2}(\mathcal{O})$ since they are piecewise smooth (they are differentiable *a.e.*, with (distributional) derivative lying in $L^2(\mathcal{O})$, which follows from equations (2) and (3) below, if $\text{Vol}(\mathcal{O}, \varphi^* \text{can}) < \infty$, which we may assume if $V_{PC}(\mathcal{O}, \mathbb{S}^n) < \infty$). Thus,

$$(4.1) \quad \lambda_1(\mathcal{O}) \int_{\mathcal{O}} |X_i \circ \varphi|^2 dv_g \leq \int_{\mathcal{O}} |\nabla X_i \circ \varphi|^2 dv_g.$$

Summing, we see that

$$(4.2) \quad \begin{aligned} \lambda_1(\mathcal{O}) \text{Vol}(\mathcal{O}, g) &= \lambda_1(\mathcal{O}) \int_{\mathcal{O}} \sum_{i=1}^{n+1} |X_i \circ \varphi|^2 dv_g \leq m \int_{\mathcal{O}} \frac{1}{m} \sum_{i=1}^{n+1} |\nabla X_i \circ \varphi|^2 dv_g \\ &\leq m \left(\int_{\mathcal{O}} \left(\frac{1}{m} \sum_{i=1}^{n+1} |\nabla X_i \circ \varphi|^2 \right)^{\frac{m}{2}} dv_g \right)^{\frac{2}{m}} \text{Vol}(\mathcal{O}, g)^{1 - \frac{2}{m}}, \end{aligned}$$

where the last inequality is Hölder's inequality.

Take a point $x \in \mathcal{O}$ at which φ is differentiable and conformal (By the definition of a PC map, almost every point of $\mathcal{O}$ satisfies these two conditions). After possibly performing an orthogonal change of coordinates at $x \in \mathcal{O}$ and on $\mathbb{R}^{n+1}$, we may assume without loss of generality that the

derivative $d\varphi(x) : T_x(\mathcal{O}) \rightarrow T_{\varphi(x)}(\mathbb{R}^{n+1})$ may be written as

$$d\varphi = \begin{bmatrix} k & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & k \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

where k is the dilatation factor of φ at x . From the fact that $\text{tr}(A^t A)$ is invariant under pre- or post-composition with an orthogonal matrix, one sees that

$$\frac{1}{m} \sum_{i=1}^{n+1} |\nabla X_i \circ \varphi|^2 = \frac{1}{m} \text{tr}(d\varphi(x)^t d\varphi(x)) = k^2.$$

Since $\varphi^*(\text{can})(x) = k^m dv_g(x)$, where can is the volume form on $\mathbb{S}^n$, we see that

$$\left(\frac{1}{m} \sum_{i=1}^{n+1} |\nabla X_i \circ \varphi|^2 \right)^{m/2} dv_g = \varphi^*(\text{can}) \text{ a.e.}$$

This yields the inequality

$$(4.3) \quad \int_{\mathcal{O}} \left(\frac{1}{m} \sum_{i=1}^{n+1} |\nabla X_i \circ \varphi|^2 \right)^{\frac{m}{2}} dv_g \leq \text{Vol}(\mathcal{O}, \varphi^* \text{can}).$$

Finally, we obtain the desired inequality by combining inequalities (4.2) and (4.3), and taking an infimum over all PC functions φ . $\square$

5. CONGRUENCE ARITHMETIC HYPERBOLIC ORBIFOLDS

This section collects some properties of arithmetic hyperbolic orbifolds which will be needed in the next section. Given a number field k , let $\mathcal{O}$ denote its ring of integers, $\mathcal{D}_k$ the absolute value of the discriminant of k and $\deg(k) = [k : \mathbb{Q}]$.

The group $H = \text{Isom}(\mathbb{H}^n)$ can be identified with $\text{O}_0(n, 1)$ — the subgroup of the orthogonal group $\text{O}(n, 1)$ which preserves the upper-half space. We now define arithmetic and congruence subgroups of H .

Definition 5.1. Let k be a totally real number field, f a quadratic form of signature $(n, 1)$ defined over k such that for every non-identity embedding $\sigma : k \rightarrow \mathbb{R}$ the form f^σ is positive definite. Then the group $\Gamma = \text{O}_0(f, \mathcal{O})$ of the integral automorphisms of f is a discrete subgroup of H via the inertia theorem. Such groups Γ and subgroups of H which are commensurable to them are called *arithmetic subgroups of the simplest type*. The field k is called a *field of definition* of Γ (and subgroups commensurable to Γ in H). An arithmetic subgroup Γ is called a *congruence subgroup* if there is a nonzero ideal $\mathfrak{a} \subset \mathcal{O}$ such that $\Gamma \supset \text{O}_0(f, \mathfrak{a})$, where $\text{O}_0(f, \mathfrak{a}) = \{g \in \text{O}_0(f, \mathcal{O}) \mid g \equiv \text{Id} \pmod{\mathfrak{a}}\}$, the *principal congruence* subgroup of $\text{O}_0(f, \mathcal{O})$.

of level $\mathfrak{a}$ (See [29] for more on arithmetic groups). We will also apply this terminology to the corresponding quotient orbifolds.

For the hyperbolic spaces of even dimension all arithmetically defined subgroups are arithmetic subgroups of the simplest type. For odd n there is another family of arithmetic subgroups given as the groups of units of appropriate Hermitian forms over quaternion algebras. Moreover, if $n = 7$ there is also the third type of arithmetic subgroups of H which are associated to the Cayley algebra. This description follows from Tits' classification of semisimple algebraic groups [37]. For our purpose it will be sufficient to consider only arithmetic subgroups of the simplest type, due to the following result of Vinberg:

Lemma 5.2. [39, Lemma 7] *If Γ contains an arithmetic subgroup generated by reflections then it is defined by a quadratic form.*

The following lemma is a special case of Proposition 3.3 in [3] which was proved as an application and extension of the method of Borel-Prasad [5].

Lemma 5.3. *Given $n > 3$, there exist positive constants c_n, δ_n such that if a maximal arithmetic subgroup $\Gamma < \text{Isom}(\mathbb{H}^n)$ is defined over the field k , then $\text{Vol}(\mathbb{H}^n/\Gamma) \geq c_n \mathcal{D}_k^{\delta_n}$.*

Proof. It was shown in [3, Sect. 3.3] that for almost all admissible fields k , if Γ is an arithmetic subgroup of $H (= \text{Isom}(\mathbb{H}^n))$ defined over k , then

$$\mu(H/\Gamma) \geq \frac{1}{2} \mathcal{D}_k^{\delta_n},$$

where μ is a Haar measure on H normalized in a certain way and δ_n is a positive constant which depends only on n and can be computed explicitly. There is a constant $c = c(n)$ such that $\text{Vol}(\mathbb{H}^n/\Gamma) = c\mu(H/\Gamma)$ (in fact, in our case c is equal to the volume of the unit sphere in $\mathbb{R}^n$). Thus for all but finitely many k , $\text{Vol}(\mathbb{H}^n/\Gamma) \geq \frac{1}{2}c\mathcal{D}_k^{\delta_n}$. We can extend this inequality to the remaining finite collection of fields by decreasing if necessary the constant c . $\square$

The next fact which we recall here may have been known for a long time. More recently, related questions in a different context were thoroughly studied by I. Horozov in his PhD thesis [16, 17]. Still we were not able to find a proof of the statement in the literature, so we will present an argument.

Lemma 5.4. *There exists a constant e_n such that if an arithmetic subgroup $\Gamma < \text{Isom}(\mathbb{H}^n)$ is defined by a quadratic form defined over k , and $C < \Gamma$ is a finite cyclic subgroup, then $|C| \leq e_n \deg(k)^{2n+2}$.*

Proof. Let Γ be a subgroup of H commensurable with the group of integral points of $G(k) = \text{O}_0(f, k)$. It is known that if the center of H is trivial (which is indeed the case for $H = \text{Isom}(\mathbb{H}^n)$) then Γ is contained in $G(k)$ [32, Prop. 1.2]. We come to a question about the torsion in $G(k) \subset \text{O}_0(f, k) \subset \text{GL}(n+1, k)$.

Let $A \in \mathrm{GL}(m, k)$, $m = n + 1$ be a torsion element of order $t \geq 2$. Let $\lambda_1, \dots, \lambda_m$ be the eigenvalues of A . As A is a matrix over k , its eigenvalues split into groups of conjugates under the action of $\mathrm{Gal}(\bar{k}/k)$. Now, $A^t = \mathrm{Id}$ implies that the eigenvalues are roots of unity. Let $t_1, \dots, t_l$ be their orders, so $t = \mathrm{lcm}(t_1, \dots, t_l)$. If λ is an eigenvalue then all its $\mathrm{Gal}(\bar{k}/k)$ -conjugates are also eigenvalues of A , which implies

$$\phi_k(t_1) + \dots + \phi_k(t_l) \leq m,$$

where $\phi_k(t)$ denotes a generalized Euler ϕ -function, which can be defined as the degree over k of the cyclotomic extension $k(\mu_t)$, where μ_t denotes a primitive t th-root of unity.

It is clear that the following inequalities are satisfied for $\phi_k(t)$ and the Euler ϕ -function:

$$\phi(t)/\deg(k) \leq \phi_k(t) \leq \phi(t).$$

To simplify the notation let $\deg(k) = d$. We have

$$\phi(t_i) \leq md.$$

This implies

$$\phi(t) \leq \phi(t_1) \dots \phi(t_l) \leq (md)^m$$

(the first inequality employs the fact that $t = \mathrm{lcm}(t_1, \dots, t_l)$).

Now using the well known inequality $\phi(t) \geq \sqrt{t}/2$, we obtain

$$t \leq 4(md)^{2m} \leq e_n d^{2n+2}.$$

□

Corollary 5.5. *Under the hypothesis of the above lemma, there exists a constant m_n such that if $F < \Gamma$ is a finite subgroup, then $|F| \leq m_n \deg(k)^{n(n+1)}$.*

Proof. By Margulis' Lemma, there is a constant M_n such that if $F < O(n)$, then there is an abelian subgroup $A < F$, such that $[F : A] \leq M_n$ (see e.g. [36, Cor. 4.1.11]). We may find common complex eigenspaces of the elements of A : $U_1, \dots, U_k$, and real eigenspaces $V_1, \dots, V_l$, where $k/2 + l \leq n$, such that A acts on U_i as a cyclic group A_i , and A acts on V_i as ± 1 . We may embed A in $\prod_{i=1}^k A_i \times (\mathbb{Z}/2\mathbb{Z})^l$, acting on $\prod_{i=1}^k U_i \times \prod_{j=1}^l V_j$. Thus $|A| \leq 2^l \prod_{i=1}^k |A_i|$. By the previous lemma, $|A_i| \leq e_n \deg(k)^{2n+2}$, since a generator of A_i is the projection of an element of A . Thus, $|F| \leq M_n |A| \leq m_n \deg(k)^{n(n+1)}$. □

Lemma 5.6. *There exists $c > 1$, such that for a number field k , $\mathcal{D}_k > c^{\deg(k)}$.*

This follows from Minkowski's theorem (see e.g. [19, V.4]), however if one wants to obtain a “good value” for the constant c , the question becomes much more complicated. There has been a considerable amount of research in this direction. We refer to [28] for a survey of the results available so far.

Lemma 5.7. *Maximal arithmetic subgroups of the simplest type are congruence.*

Proof. Let Γ be a maximal arithmetic subgroup which is commensurable with $G(\mathcal{O}) = \mathrm{O}_0(f, \mathcal{O})$ for some quadratic form f defined over k . There exists an $\mathcal{O}$ -lattice L in $V = k^{n+1}$ such that $\Gamma \cap G(k) (= \Gamma) = G^L = \{g \in G(k) \mid g(L) = L\}$ (see [29, Prop. 4.2] for $k = \mathbb{Q}$, the general case is entirely similar as we do not claim that L is a free $\mathcal{O}$ -lattice). Let also M be the standard lattice in V so that $G^M = G(\mathcal{O})$. We claim that there exists a principal congruence subgroup of G^M which is contained in G^L . The argument is the same as in [29, Lemma 4.1.1]: Let $\phi : G \rightarrow G$ be a k -isomorphism defined by the change of basis of V from the standard basis (of M) to a k -basis contained in L . It follows that if Δ is a principal congruence subgroup of G^M with respect to an ideal $\mathfrak{a} \subset \mathcal{O}$ such that the coefficients of $\mathfrak{a}(\phi - \mathrm{Id})$ in the standard basis are integral, then $\Delta \subset G^L$. $\square$

Remark: The results which were discussed in this section so far are valid for arbitrary arithmetic subgroups, the proofs for the general case are similar to the case of the simplest type which we considered in detail.

The following theorem was stated in the case of arithmetic groups defined over $\mathbb{Q}$, but is true in general. It is a translation of results of Jacquet-Langlands [18] and Gelbart-Jacquet [13] in the theory of automorphic forms into a statement about eigenvalues of congruence arithmetic groups, and it generalizes a famous theorem of Selberg for congruence subgroups of $\mathrm{PSL}(2, \mathbb{Z})$ [34].

Theorem 5.8 (Vigneras [38]). *Let $\mathcal{O} = \mathbb{H}^2/\Gamma$, where Γ is an arithmetic congruence Fuchsian subgroup of $\mathrm{Isom}(\mathbb{H}^2)$. Then $\lambda_1(\mathcal{O}) \geq \frac{3}{16}$.*

In the above theorem, it is conjectured that $\lambda_1(\mathcal{O}) \geq \frac{1}{4}$, which generalizes a conjecture of Selberg about congruence subgroups of $\mathrm{PSL}(2, \mathbb{Z})$. The following theorem of Burger and Sarnak extends this result to higher dimensions, by inducing from the 2-dimensional case.

Theorem 5.9. [8, Cor. 1.3(a)] *Let $\mathcal{O} = \mathbb{H}^n/\Gamma$, where Γ is an arithmetic congruence subgroup defined by a quadratic form, $n \geq 3$ (or if $n = 3$, Γ need only be arithmetic congruence). Then $\lambda_1(\mathcal{O}) \geq \frac{2n-3}{4}$.*

Under the hypotheses of Theorem 5.9, it is conjectured that $\lambda_1(\mathcal{O}) \geq n-2$, which is a special case of the *generalized Ramanujan conjecture* [8, Cor. 1.3(b)].

6. FINITENESS OF ARITHMETIC HYPERBOLIC MAXIMAL REFLECTION GROUPS

We put together the results in the previous sections to prove our main theorem.

Theorem 6.1. *There are only finitely many conjugacy classes of arithmetic maximal reflection groups in $\mathrm{Isom}(\mathbb{H}^n)$.*

Proof. Suppose that Γ is an arithmetic maximal reflection group. Then there is no group $\Gamma' < \text{Isom}(\mathbb{H}^n)$ such that $\Gamma < \Gamma'$ and Γ' is generated by reflections. There does exist a maximal lattice $\Gamma_0 < \text{Isom}(\mathbb{H}^n)$ satisfying $\Gamma \leq \Gamma_0$. The group Γ is generated by reflections in a finite volume polyhedron P , which forms a fundamental domain for Γ .

Lemma 6.2 (Vinberg [39]). *Γ is a normal subgroup of Γ_0 . Moreover, there is a finite subgroup $\Theta < \Gamma_0$ such that $\Theta \rightarrow \Gamma_0/\Gamma$ is an isomorphism, and Θ is the group of symmetries of the polyhedron P .*

Proof. This follows from the fact that the set of reflections in Γ_0 is conjugacy invariant, and therefore the subgroup of Γ_0 generated by reflections in Γ_0 is normal in Γ_0 . Since Γ is a maximal reflection group, this subgroup of Γ_0 must be Γ .

Suppose there is an element $\gamma \in \Gamma_0$ such that $\text{int}(P) \cap \gamma(\text{int}(P)) \neq \emptyset$ and $\gamma(P) \neq P$. Then there is a geodesic plane V containing a face of P such that $\gamma(V) \cap \text{int}(P) \neq \emptyset$. The reflection $r_V \in \Gamma$ in the plane V is conjugate to a reflection $r_{\gamma(V)} = \gamma r_V \gamma^{-1}$, which is not in Γ since $r_{\gamma(V)}(\text{int}(P)) \cap \text{int}(P) \neq \emptyset$. However $\Gamma \trianglelefteq \Gamma_0$ implies $r_{\gamma(V)} \in \Gamma$, a contradiction.

Let Θ be the subgroup of Γ_0 such that $\Theta(P) = P$. Θ is finite because P is finite volume and has finitely many faces.

Consider the composition $\Theta \hookrightarrow \Gamma_0 \rightarrow \Gamma_0/\Gamma$. $\Gamma \cap \Theta$ is trivial because P is a fundamental domain for Γ . The composition map is therefore injective. To see surjectivity, pick a $\gamma_0 \in \Gamma_0$ and find a $\gamma \in \Gamma$ such that $\gamma_0^{-1}(\text{int}(P)) \cap \gamma(\text{int}(P)) \neq \emptyset$. Then $\gamma_0\gamma(P) = P$, so $\gamma_0\gamma = \theta \in \Theta$. Therefore $\gamma_0\Gamma = \theta\Gamma$. This shows that the composition $\Theta \rightarrow \Gamma_0/\Gamma$ is surjective, and an isomorphism. $\square$

Let $\mathcal{O} = \mathbb{H}^n/\Gamma_0$, and let Θ be the finite group coming from the previous lemma.

Now, we remark that the case of $n = 2$ follows from the result of Long-MacLachlan-Reid [21]. If $n = 2$ then there is an index 2 subgroup $\Gamma' < \Gamma_0$ which is a genus 0 Fuchsian group. Thus, by Theorem 1.1 of [21], we can conclude that there are only finitely many possible Γ' , since Γ' is also a congruence group, and therefore there are only finitely many Γ_0 and Γ . In fact, $\text{Area}(\mathbb{H}^2/\Gamma') < \frac{128}{3}\pi$. Also, the case of $n = 3$ was proven in [1]. So we may restrict to the case that $4 \leq n \leq 995$, by Prokhorov's Theorem [30]. In fact, since Nikulin has shown the finiteness of maximal non-uniform arithmetic reflection groups [24, 25, 26], we may restrict to the case that Γ is cocompact, and therefore $n < 30$, although our argument gives a new proof of Nikulin's Theorem, making use of Prokhorov's Theorem.

Consider $\mathbb{H}^n \subset \mathbb{S}^n$ embedded conformally as the upper hemisphere of $\mathbb{S}^n$, so that $\text{Isom}(\mathbb{H}^n)$ acts conformally on $\mathbb{S}^n$. Normalize so that Θ acts isometrically on $\mathbb{S}^n$, by placing the common fixed point of Θ at the north pole of $\mathbb{S}^n$. Clearly, $V_{PC}(n, \mathcal{O}) = V_{PC}(n, P/\Theta)$, since $|\mathcal{O}| = |P/\Theta|$. Then the orbifold $P/\Theta \subset \mathbb{S}^n/\Theta$ is a conformal embedding, so fact 4 of section 2 gives the inequality $V_{PC}(n, P/\Theta) \leq V_{PC}(n, \mathbb{S}^n/\Theta)$.

We now apply the eigenvalue estimates from Theorems 4.2 and 5.9:

$$\frac{2n-3}{4} \text{Vol}(\mathcal{O})^{\frac{2}{n}} \leq \lambda_1(\mathcal{O}) \text{Vol}(\mathcal{O})^{\frac{2}{n}} \leq n V_{PC}(n, \mathcal{O})^{\frac{2}{n}} \leq n (V_{PC}(\mathbb{S}^n / \Theta))^{\frac{2}{n}}.$$

Assume now that Γ_0 is defined over the number field k . By Theorem 3.1 and Corollary 5.5 we have

$$V_{PC}(\mathbb{S}^n / \Theta) \leq C(n) |\Theta|^{2n} \leq C(n) (m_n \deg(k)^{2(n+1)n})^{2n}.$$

By Lemmas 5.3 and 5.6, we have $\text{Vol}(\mathcal{O}) = \text{Vol}(\mathbb{H}^n / \Gamma_0) \geq c_n c^{\delta_n \deg(k)}$. Putting these inequalities together, we obtain an inequality of the form

$$c^{\delta_n \deg(k)} \leq C_1(n) \deg(k)^{4(n+1)n^2},$$

which implies that $\deg(k)$ must be bounded.

We also obtain

$$\text{Vol}(\mathcal{O})^{\frac{2}{n}} \leq \frac{4n}{2n-3} C(n)^{\frac{2}{n}} m_n^4 \deg(k)^{8(n+1)n}.$$

Since $\deg(k)$ is bounded, this gives an upper bound on $\text{Vol}(\mathcal{O})$ depending only on n .

Since volumes of arithmetic hyperbolic orbifolds are discrete [4, 41], and Γ is uniquely determined by Γ_0 , we conclude that there are only finitely many arithmetic maximal reflection groups in dimension n . $\square$

7. CONCLUSION

It is clear that for a finite volume polyhedron $P \subset \mathbb{H}^n$, $V_{PC}(n, P) = \text{Vol}(\mathbb{S}^n)$, so by Theorem 4.2

$$\lambda_1(P) \text{Vol}(P)^{\frac{2}{n}} \leq n \text{Vol}(\mathbb{S}^n)^{\frac{2}{n}}.$$

Thus in n dimensions, there are finitely many reflection groups $\Gamma < \text{Isom}(\mathbb{H}^n)$ which have a lower bound on $\lambda_1(\mathbb{H}^n / \Gamma)$. If it were true that maximal arithmetic hyperbolic reflection groups are congruence, then this would simplify the argument in this paper quite a bit, since we would immediately get an upper bound on the volume by the above observation. So it is natural to ask, does there exist a non-congruence maximal arithmetic reflection group?

The following conjecture seems to be interesting, and would also give an alternative argument to the main theorem of this paper:

Conjecture 7.1. There is a function $K(n)$, such that if $\mathcal{O}$ is an elliptic n -orbifold, then $V_{PC}(\mathcal{O}) \leq K(n)$.

Another approach to the proof of the main theorem would be possible if we assume the validity of the *short geodesic conjecture* for arithmetic hyperbolic n -orbifolds. As it was pointed out earlier in this section, Theorem 4.2 implies

$$\lambda_1(P) \text{Vol}(P)^{\frac{2}{n}} \leq n \text{Vol}(\mathbb{S}^n)^{\frac{2}{n}}.$$

Now if the group Γ generated by reflections in P is arithmetic then it is contained in some maximal arithmetic subgroup $\Gamma_0 < \text{Isom}(\mathbb{H}^n)$ with a finite index l . Let k be the field of definition of Γ_0 . The corollary from Lemma 5.4

and Lemma 6.2 imply that $l \leq m_n \deg(k)^{2(n+1)^n}$, and Lemmas 5.3, 5.6 imply that $\text{Vol}(P) \geq l c^{\delta_n \deg(k)}$. What remains is to estimate $\lambda_1(P) = \lambda_1(\Gamma)$ using the fact that $\lambda_1(\Gamma_0) \geq (2n-3)/4$ which follows from the Theorem 5.9. This brings us to the question about the spectrum of coverings which was studied by Brooks in [6, 7]. The result of [7] implies that $\lambda_1(\Gamma) \geq c(\Gamma_0) \lambda_1(\Gamma_0)/l^2$ and more careful analysis of the proof shows that the constant $c(\Gamma_0)$ depends only on the lower bound for the injectivity radius of $\mathbb{H}^n/\Gamma_0$. The short geodesic conjecture says that in every dimension $n \geq 3$ there exist a universal lower bound for the injectivity radius, so assuming the conjecture is true we can write $\lambda_1(\Gamma) \geq c(n) \lambda_1(\Gamma_0)/l^2$. Substituting this to the main inequality together with the bounds for the index l and $\text{Vol}(P)$ we are getting the finiteness results in a similar way to the proof of the theorem in sect. 6. This argument avoids the analysis of the conformal volumes of spherical orbifolds (sect. 3) but unfortunately is based on an open conjecture.

From the main theorem, we conclude that given an arithmetic reflection group $\Gamma < \text{Isom}(\mathbb{H}^n)$, it must lie in one of finitely many maximal reflection groups (up to conjugacy). If Γ is a reflection group in a polyhedron P for which all the dihedral angles are $\pi/2$ or all are $\pi/3$, then there are infinitely many finite covolume reflection subgroups of Γ . Thus we see that there are commensurability classes of arithmetic groups for which there are infinitely many reflection groups in the commensurability class, and thus in our finiteness result, the maximality assumption is crucial. The same is true for the arithmeticity assumption. For example, consider the groups generated by reflections of the hyperbolic plane in the sides of a hyperbolic triangle with angles $\pi/2, \pi/3, \pi/m$ ($m \geq 7$). These groups are known to be maximal discrete subgroups of $\text{Isom}(\mathbb{H}^2)$ but all except finitely many of them are non-arithmetic. A similar construction is available for hyperbolic 3-space with triangular prisms replacing triangles [22].

It is an interesting project to try to identify all arithmetic maximal reflection groups and to classify their reflection subgroups of finite covolume. Although all the constants in our finiteness argument are computable, the quantitative estimates which could be deduced directly from the proof are unreasonably large. However, if we restrict to the congruence reflection groups (see the discussion in the beginning of this section) or make some additional assumptions on the groups of automorphisms of the fundamental polyhedra of the reflection groups, then the bounds can be improved essentially.

REFERENCES

- [1] I. Agol, *Finiteness of arithmetic Kleinian reflection groups*, Proceedings of ICM 2006, Madrid, vol. II, European Mathematical Society, Zurich, 2006, arXiv:math.GT/0512560, pp. 951–960.
- [2] D. Allcock, *Infinitely many hyperbolic Coxeter groups through dimension 19*, Geom. Topol. **10** (2006), 737–758 (electronic).
- [3] M. Belolipetsky, *Counting maximal arithmetic subgroups*, preprint, arxiv:math.GR/0501198.

- [4] A. Borel, *Commensurability classes and volumes of hyperbolic 3-manifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **8** (1981), no. 1, 1–33.
- [5] A. Borel and G. Prasad, *Finiteness theorems for discrete subgroups of bounded covolume in semi-simple groups*, Inst. Hautes Études Sci. Publ. Math. (1989), no. 69, 119–171.
- [6] R. Brooks, *The fundamental group and the spectrum of the Laplacian*, Comment. Math. Helv. **56** (1981), no. 4, 581–598.
- [7] ———, *The spectral geometry of a tower of coverings*, J. Differential Geom. **23** (1986), no. 1, 97–107.
- [8] M. Burger and P. Sarnak, *Ramanujan duals. II*, Invent. Math. **106** (1991), no. 1, 1–11.
- [9] J. B. Dennin, Jr., *Fields of modular functions of genus 0*, Illinois J. Math. **15** (1971), 442–455.
- [10] ———, *Subfields of $K(2^n)$ of genus 0*, Illinois J. Math. **16** (1972), 502–518.
- [11] A. El Soufi and S. Ilias, *Immersiones minimales, première valeur propre du laplacien et volume conforme*, Math. Ann. **275** (1986), no. 2, 257–267.
- [12] J. Elstrodt, F. Grunewald, and J. Mennicke, *Groups acting on hyperbolic space*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 1998, Harmonic analysis and number theory.
- [13] S. Gelbart and H. Jacquet, *A relation between automorphic representations of $GL(2)$ and $GL(3)$* , Ann. Sci. École Norm. Sup. (4) **11** (1978), no. 4, 471–542.
- [14] M. L. Gromov and V. A. Rohlin, *Imbeddings and immersions in Riemannian geometry*, Uspehi Mat. Nauk **25** (1970), no. 5 (155), 3–62.
- [15] J. Hersch, *Quatre propriétés isopérimétriques de membranes sphériques homogènes*, C. R. Acad. Sci. Paris Sér. A-B **270** (1970), A1645–A1648.
- [16] I. Horozov, *Euler characteristics of arithmetic groups*, Ph.D. thesis, Brown University, 2003, arXiv:math.GR/0311117.
- [17] ———, *Euler characteristics of arithmetic groups*, Math. Res. Lett. **12** (2005), no. 2-3, 275–291.
- [18] H. Jacquet and R. P. Langlands, *Automorphic forms on $GL(2)$* , Springer-Verlag, Berlin, 1970, Lecture Notes in Mathematics, Vol. 114.
- [19] S. Lang, *Algebraic number theory*, Addison-Wesley Publishing Co., Inc., Reading, Mass.-London-Don Mills, Ont., 1970.
- [20] P. Li and S. T. Yau, *A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces*, Invent. Math. **69** (1982), no. 2, 269–291.
- [21] D. D. Long, C. Maclachlan, and A. W. Reid, *Arithmetic Fuchsian groups of genus zero*, Pure Appl. Math. Q. **2** (2006), no. 2, 569–599.
- [22] Colin Maclachlan and Alan W. Reid, *The arithmetic of hyperbolic 3-manifolds*, Graduate Texts in Mathematics, vol. 219, Springer-Verlag, New York, 2003.
- [23] G. A. Margulis, *Discrete groups of motions of manifolds of nonpositive curvature*, Proceedings of the International Congress of Mathematicians (Vancouver, B.C., 1974), Vol. 2, Canad. Math. Congress, Montreal, Que., 1975, pp. 21–34.
- [24] V. V. Nikulin, *On the arithmetic groups generated by reflections in Lobachevskii spaces*, Izv. Akad. Nauk SSSR Ser. Mat. **44** (1980), no. 3, 637–669, 719–720.
- [25] ———, *On the classification of arithmetic groups generated by reflections in Lobachevskii spaces*, Izv. Akad. Nauk SSSR Ser. Mat. **45** (1981), no. 1, 113–142, 240.
- [26] ———, *Discrete reflection groups in Lobachevsky spaces and algebraic surfaces*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986) (Providence, RI), Amer. Math. Soc., 1987, pp. 654–671.
- [27] V. V. Nikulin, *Finiteness of the number of arithmetic groups generated by reflections in Lobachevsky spaces*, preprint 2006, arXiv:math.AG/0609256, 5 pages.
- [28] A. M. Odlyzko, *Bounds for discriminants and related estimates for class numbers, regulators and zeros of zeta functions: a survey of recent results*, Sémin. Théor. Nombres Bordeaux (2) **2** (1990), no. 1, 119–141.
- [29] V. Platonov and A. Rapinchuk, *Algebraic groups and number theory*, Pure and Applied Mathematics, vol. 139, Academic Press Inc., Boston, MA, 1994, Translated from the 1991 Russian original by Rachel Rowen.
- [30] M. N. Prokhorov, *Absence of discrete groups of reflections with a noncompact fundamental polyhedron of finite volume in a Lobachevskii space of high dimension*, Izv. Akad. Nauk SSSR Ser. Mat. **50** (1986), no. 2, 413–424.

- [31] M. Reed and B. Simon, *Methods of modern mathematical physics. IV. Analysis of operators*, Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1978.
- [32] A. A. Ryzhkov and V. I. Chernousov, *On the classification of maximal arithmetic subgroups of simply connected groups*, Mat. Sb. **188** (1997), no. 9, 127–156.
- [33] P. Scott, *Subgroups of surface groups are almost geometric*, Journal of the London Mathematical Society. Second Series **17** (1978), no. 3, 355–565.
- [34] A. Selberg, *On the estimation of Fourier coefficients of modular forms*, Proc. Sympos. Pure Math., Vol. VIII, Amer. Math. Soc., Providence, R.I., 1965, pp. 1–15.
- [35] G. Szegő, *Inequalities for certain eigenvalues of a membrane of given area*, J. Rational Mech. Anal. **3** (1954), 343–356.
- [36] W. P. Thurston, *Three-dimensional geometry and topology. Vol. 1*, Princeton Mathematical Series, vol. 35, Princeton University Press, Princeton, NJ, 1997, Edited by Silvio Levy.
- [37] J. Tits, *Classification of algebraic semisimple groups*, Algebraic Groups and Discontinuous Subgroups (Proc. Sympos. Pure Math., Boulder, Colo., 1965), Amer. Math. Soc., Providence, R.I., 1966, 1966, pp. 33–62.
- [38] M.-F. Vignéras, *Quelques remarques sur la conjecture $\lambda_1 \geq \frac{1}{4}$* , Seminar on number theory, Paris 1981–82 (Paris, 1981/1982), Progr. Math., vol. 38, Birkhäuser Boston, Boston, MA, 1983, pp. 321–343.
- [39] È. B. Vinberg, *Discrete groups generated by reflections in Lobachevskii spaces*, Mat. Sb. (N.S.) **72** (114) (1967), 471–488; correction, ibid. 73 (115) (1967), 303.
- [40] ———, *Absence of crystallographic groups of reflections in Lobachevskii spaces of large dimension*, Trudy Moskov. Mat. Obshch. **47** (1984), 68–102, 246.
- [41] H. C. Wang, *Topics on totally discontinuous groups*, Symmetric spaces (Short Courses, Washington Univ., St. Louis, Mo., 1969–1970), Dekker, New York, 1972, pp. 459–487. Pure and Appl. Math., Vol. 8.
- [42] P. C. Yang and S. T. Yau, *Eigenvalues of the Laplacian of compact Riemann surfaces and minimal submanifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **7** (1980), no. 1, 55–63.
- [43] P. Zograf, *A spectral proof of Rademacher’s conjecture for congruence subgroups of the modular group*, J. Reine Angew. Math. **414** (1991), 113–116.

MSCS UIC 322 SEO, M/C 249, 851 S. MORGAN ST., CHICAGO, IL 60607-7045, USA
E-mail address: `agol@math.uic.edu`

DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY, SOUTH RD, DURHAM, DH1 3LE, UK
E-mail address: `mikhail.belolipetsky@durham.ac.uk`

STANFORD UNIVERSITY, MATHEMATICS, BLDG. 380, 450 SERRA MALL, STANFORD, CA 94305-2125, USA
E-mail address: `storm@math.stanford.edu`

MSCS UIC 322 SEO, M/C 249, 851 S. MORGAN ST., CHICAGO, IL 60607-7045, USA
E-mail address: `kwhyte@math.uic.edu`